

Sleep and Recovery in Team Sport: Current Sleep-Related Issues Facing Professional Team-Sport Athletes

Hugh H.K. Fullagar, Rob Duffield, Sabrina Skorski, Aaron J. Coutts, Ross Julian, and Tim Meyer

While the effects of sleep loss on performance have previously been reviewed, the effects of disturbed sleep on recovery after exercise are less reported. Specifically, the interaction between sleep and physiological and psychological recovery in team-sport athletes is not well understood. Accordingly, the aim of the current review was to examine the current evidence on the potential role sleep may play in postexercise recovery, with a tailored focus on professional team-sport athletes. Recent studies show that team-sport athletes are at high risk of poor sleep during and after competition. Although limited published data are available, these athletes also appear particularly susceptible to reductions in both sleep quality and sleep duration after night competition and periods of heavy training. However, studies examining the relationship between sleep and recovery in such situations are lacking. Indeed, further observational sleep studies in team-sport athletes are required to confirm these concerns. Naps, sleep extension, and sleep-hygiene practices appear advantageous to performance; however, future proof-of-concept studies are now required to determine the efficacy of these interventions on postexercise recovery. Moreover, more research is required to understand how sleep interacts with numerous recovery responses in team-sport environments. This is pertinent given the regularity with which these teams encounter challenging scenarios during the course of a season. Therefore, this review examines the factors that compromise sleep during a season and after competition and discusses strategies that may help improve sleep in team-sport athletes.

Keywords: regeneration, exercise, stress, soccer, circadian rhythms

High-performance team-sport athletes endure numerous physiological, psychological, and neuromuscular stressors during training and competition.¹ It is logical that these athletes balance these stressors with appropriate recovery to maximize performance and adaptation, while also minimizing injury risk.² A crucial part of this stress-recovery balance is the management of an athlete's sleep, especially during intense training and competition.³ However, while the interest afforded to the relationship between sleep and athletic performance is well documented,⁴ the evidence underpinning the role of sleep in recovery is less understood. This is surprising from both a scientific and an applied perspective, given that athletes often rate sleep as their most important recovery strategy.⁵

There are 3 key factors that determine the recuperative outcome of sleep: the duration (total sleep time), quality, and phase (circadian timing) of sleep.⁶ A "healthy" night of sleep has been suggested to be 7 to 9 hours.⁷ In addition to duration, sleep quality is also critical for optimal health and restorative functioning.⁷ Although a clear definition is not readily available, sleep quality can best be described as the personal satisfaction with the sleep experience.⁷ Furthermore, the timing of sleep will also influence the effectiveness of the sleep bout. The timing of an individual's preferred bedtime in turn affects his or her circadian rhythms (ie, body temperature, hormone regulation), which can affect both sleep duration and sleep quality.⁶ From an athletic perspective, disturbances to 1 or

all of these collective aspects of sleep are suggested to affect the postexercise recovery process.⁶ For instance, it has been shown that a reduction in the quantity and quality of sleep hinders the capacity of rugby league footballers to recover for the demands of ensuing training and competitive bouts.⁸ Thus, it may be paramount for team-sport athletes to be aware of situations where disturbed sleep duration, quality, or phase may affect ensuing recovery.

A reduction in sleep duration and/or quality in individual athletes before⁹⁻¹¹ and during competition¹² has been recently documented. While there is less information available on team-sport athletes, Lastella et al¹³ reported a mean sleep duration of 7.0 h/night in 58 elite Australian team-sport athletes during a typical training phase, ~1 hour less than the recommended 8 h/night. Further to these findings, sleep disruption or deprivation can occur for team-sport athletes, particularly during short- or long-haul travel,¹⁴⁻¹⁶ congested competition schedules,¹ and training or playing at night,¹⁷ presenting the potential for compromised recovery.^{3,8} Indeed, sleep loss in team-sport athletes is often affected by these situational factors,¹⁸ with many professional teams currently facing the challenge of coping with these specific but recurring stressors. For example, Major League Baseball players play every 2 days combined with repeated travel across the United States, which provides conditions that are not conducive to optimal sleep.¹⁹ Similarly, the majority of European soccer tournaments are commonly played at night, resulting in late-night finishes and players subjectively reporting sleep loss.²⁰ These observations of altered sleep in team-sport athletes are also supported by objective evidence of postcompetitive sleep disturbance in elite rugby union players¹⁷ and professional Australian soccer players.¹⁶ Furthermore, a recent report that 52.3% of elite (individual and team-sport) athletes experience sleep disturbances after late matches or training sessions.¹⁸ Collectively, these data suggest that although "normal" sleep patterns may be sufficient,

Fullagar, Skorski, Julian, and Meyer are with the Inst of Sport and Preventive Medicine, Saarland University, Saarbrücken, Germany. Duffield and Coutts are with the Sport & Exercise Discipline Group, UTS: Health, University of Technology, Sydney, Australia. Address author correspondence to Hugh Fullagar at hugh.fullagar@uni-saarland.de.

under specific, recurring circumstances there are cases for reduced sleep duration and quality in team-sport athletes.

At present, the importance of sleep as a recovery method in team-sport athletes (ie, return to baseline of psychophysiological and performance parameters after exercise and disrupted sleep) is unclear. In particular, there is little analysis of the role sleep plays in the postexercise recovery process during various situations where sleep is compromised. While the literature examining the interaction between sleep and recovery in athletes is increasing (Figure 1), there have been no critical reviews of these factors in the context of training and competition demands of team-sport athletes. Accordingly, the aim of the current study was to examine the evidence of the potential role sleep may play in postexercise recovery, with a specific focus on professional team-sport athletes. As such, an analysis of situations that may continually compromise sleep throughout a season and/or one-off postcompetition sleep disturbance is provided. Strategies to alleviate such issues facing team-sport athletes are also addressed. For this review, it is important to discern the difference between recovery and performance. From an athletic perspective, performance in absolute terms refers to the context and magnitude to which athletes complete certain tasks in their sporting domain.²¹ These can include but are not limited to competition performance (eg, goals scored by a footballer), predictors of performance (eg, sprinting speed), and surrogate measures of performance (eg, countermovement-jump score). The effects of sleep loss on performance trials involve baseline performance measures followed by a sleep-loss intervention/sleep-control condition and then final performance measures the next morning. Comparatively, recovery refers to the degree at which parameters return to baseline after a distinct exercise bout and disrupted sleep (eg, return of creatine kinase to baseline values after a rugby match or the return of YoYo

test performance to baseline values after a training session).^{6,8} Thus, the main discernible difference between performance and recovery is that recovery experiments follow a distinct time-course analysis from a prior stressor (ie, match play). This makes them suitable for the assessment of the health, well-being, and readiness to perform of team-sport athletes.

Sleep and Recovery for Team-Sport Athletes

A typical night of sleep is composed of approximately 90-minute cycles divided into periods of rapid-eye-movement (REM) and non-REM (NREM) sleep. While REM sleep has a role in periodic brain activation, localized recuperative processes, and emotional regulation, the role for NREM sleep is proposed to assist with energy conservation and nervous system recuperation.²² Taken collectively, there is considerable evidence supporting the recuperative nature of sleep in restoring molecular homeostasis, cellular maintenance, and synaptic plasticity.^{6,22,23} From an athletic perspective, this implies that disturbances to either the timing of sleep phases or the quality and duration of sleep within these phases can result in the hindrance of psychological and physical recovery after an exercise bout.⁶ This would seem especially pertinent for field-based team-sport athletes who are typically exposed to prolonged bouts of intermittent-sprint activity during both high-intensity training and competition. Logically, exposure to such activity will increase the need for recovery and subsequently increase the overall requirement for sleep.¹³

From this perspective, it seems rational to first investigate the sleep-wake behavior of team-sport athletes during and after training and competition periods. Mah et al²⁴ reported mean sleep durations




Figure 1 — The increase in the number of sleep, athlete, and recovery publications over the past 8 years. The solid fill lines illustrate the amount of literature that appears in a Pub Med database search using the terms *sleep*, *recovery*, and *athlete* in all fields for each calendar year.

of 6.7 ± 1.0 hours in college basketballers during a competitive season. Similarly, Lastella et al¹³ found that a sample of 58 elite Australian team-sport athletes slept for a mean duration of 7.0 ± 1.2 hours during a regular training phase. With regard to sleep following competition, Eagles et al¹⁷ found a significant reduction in sleep duration on game nights compared with nongame nights.¹⁷ Juliff et al¹⁸ reported that more than half of a sample of 283 elite individual and team-sport athletes (of which 210 were from team sports) endured sleep disturbances after a late training session or match.¹⁸ In support of this, sleep duration and quality were significantly reduced on the night of away matches compared with the night prior in elite Australian soccer players.¹⁶ While caution needs to be taken in comparing these studies (ie, due to differences in sleep-assessment methodologies), it seems reasonable to assume that sleep in team-sport athletes depends on many factors. These could include the type of sport, training demands, age, time of season, and team culture.¹³ Overall, high-performance team-sport athletes are considered susceptible to sleep loss during training periods and after match play (especially after night matches). While such insight is important, further descriptive research of sleep with high-performance team-sport athletes is required to confirm this, most importantly for the nights after competition.

Recent studies have also shown that sleep restriction after team-sport competition affects the time course of recovery for both performance and psychophysiological measures. For instance, Skein et al⁸ investigated the effect of sleep deprivation (0 h sleep) compared with normal sleep (~8 h) on the physiological and perceptual recovery of 11 rugby league footballers after competitive matches in a randomized crossover design. Overall, sleep deprivation negatively affected recovery, with significant impairments observed in mean and peak countermovement-jump height and cognitive reaction time. Although sleep deprivation was excessive, this study highlights the increased physiological load during wakefulness after sleep loss in team sports and, in turn, suppression of cognitive function and lower-body power. Similarly, Fowler et al¹⁶ reported significant reductions in sleep duration and quality, along with an impaired stress-recovery balance, on the night of a match compared with the night prior for away matches. While additional literature is lacking

in team-sport athletes, there is further evidence of this relationship in individual athletes. For instance, significant reductions in sleep quantity and efficiency were associated with increased fatigue and impaired exercise capacity in a group of 10 functionally overreached elite synchronized swimmers.²⁵ Furthermore, McMurray and Brown²⁶ investigated the cardiovascular and metabolic responses of 5 participants during submaximal exercise after 24 hours of sleep deprivation. They reported increased minute ventilation and oxygen uptake during the recovery period, suggesting negative effects of sleep loss on physiological recovery.²⁶ Nonetheless, the evidence as to how sleep interacts with multifactorial recovery responses in high-performance team-sport environments is currently lacking. In particular, there are few data on longitudinal objective sleep available in the scientific literature. This is surprising given that this would appear the first step in understanding the relationship between sleep and recovery.

Finally, since a variety of other recovery strategies are used in sport, some studies have also examined the interaction between sleep and these protocols. For instance, Robey et al²⁷ reported that cold-water immersion posttraining does not affect subsequent sleep duration, onset, or efficiency. However, the mechanisms between the interaction of sleep and other recovery protocols are difficult to determine, due to an abundance of confounding factors (eg, protocol type, timing, facilities). Further research and practical investigation in professional environments that address whether it is more advantageous to use a recovery protocol that enhances sleep and/or whether a combination of these protocols enhances the recovery process are warranted. This is especially pertinent given the wide prevalence of these methods in team sports.

Sleep-Related Issues Facing Team-Sport Athletes

As summarized in Figure 2, the following section outlines particular situations where sleep is at risk of compromise in team-sport athletes. While acknowledging the previous work done in this area but also recognizing the absence of published data over prolonged




Figure 2 — A schematic representation of the commonly encountered situations in team sports that may compromise sleep patterns and potentially recovery. Theoretical effects of these situations are also described; however, it should be noted that more research is required to confirm the majority of these effects.

periods, this section gives particular relevance to situations during a season and/or one-off postcompetition sleep disturbance.

Team-Sport Matches Played at Night

As often determined by television scheduling, numerous team sports schedule their matches at night. Indeed, the pure timing of matches (ie, some matches in the Spanish La Liga commence at 10:00 PM) will force players into later bedtimes.¹ Furthermore, since physical activity promotes arousal, it has long been assumed that exercising during the evening hours produces a greater number of sleep disturbances than exercising during daylight.²⁰ Team-sport athletes also have extensive postgame commitments such as press conferences, recovery practices, and social functions, which could lead to later bedtimes and disrupt sleep duration and quality.¹ As alluded to previously, Juliff et al¹⁸ found that 52.3% of a sample of 283 elite individual ($n = 73$) and team-sport ($n = 210$) athletes reported sleep disturbances after a night training session or match. Moreover, 59.1% of team-sport athletes reported that they did not use a strategy to overcome these sleep disturbances.¹⁸ Furthermore, a recent review on regenerative interventions used in professional soccer explained that many medical doctors report that players lose sleep after night matches, including findings on elite Bundesliga soccer players subjectively reporting reduced sleep duration and quality.²⁰ Notwithstanding these findings, the anecdotal evidence of athletes reporting sleep disturbances after night competition outweighs that documented in the literature; thus, further research in elite athletic populations is required to confirm this.

Recent data show that performing maximal aerobic exercise in the evening results in elevated sleep-onset latency, awakenings, and REM-sleep latency—suggesting poorer overall sleep quality in judo competitors.²⁸ While several physiological variables are elevated before sleep onset after late-night vigorous exercise (suggesting possible effects on cardiac autonomic control and metabolic function²⁹), delayed sleep onset can also be caused by mental stimulation or cognitive fatigue.²³ Moreover, given that pain is a significant predictor of a poor night's sleep,³⁰ it is likely that prolonged late-night, high-intensity exercise (equivalent to match situations) will incur sleep disturbances throughout the night as a result of pain and soreness. This is of particular relevance for heavy-contact sports such as American football, ice hockey, and rugby union. It should be noted that there is opposing evidence on the effect of competing at night on sleep. For instance, Roach et al³¹ reported no effect of 2 night (7:00 to 9:00 PM) matches on sleep in elite junior soccer players. Similarly, Robey et al³² found no effect of early-evening high-intensity training on the subsequent sleep quality or duration in elite youth soccer players.

In light of this, it should be recognized that the mechanisms behind the effects of exercise (and its timing) on sleep are complex due to the main confounding variable (among others) of the stress induced by the exercise itself. From an applied perspective, future research must first focus on providing objective evidence (eg, acute and chronic measurements of ActiGraphy) on whether disturbances after match play at night occur. Researchers might also focus on the effects of disrupted sleep after match play in team-sport athletes and attempt to delineate the mechanisms responsible. At present, practitioners should also be aware of the intraindividual variability in sleep requirement and chronotype (those who rise early in the morning vs those who prefer later bedtimes). Accommodating these differences within a team environment is difficult as it may require more individualized approaches. Indeed, this would be even more pertinent for teams scheduling training the day after a game. For

instance, training in the absence of sufficient sleep after late-night matches may potentiate negative outcomes. This may create recovery concerns given that players will sleep differently after these matches, while also possibly placing those who are training at an unnecessary risk of injury.³³

Sleep and Travel Fatigue

Cumulative sleep loss occurs as a consequence of travel during busy periods, which tends to lead to cumulative fatigue over a season.³⁴ Travel fatigue is dependent on the distance and frequency of travel and the length of the season. It should be noted that travel-induced fatigue is separate from jet-lag fatigue, with the main difference being that jet-lag comprises an effect of time-zone change.³⁴ The influences of jet-lag arising from long-haul international travel in elite athletes have been discussed previously^{34,35} and thus will not be further addressed here. Sleep disturbances during or after travel can result in reductions in mood, acute fatigue, and difficulty in initiating sleep at the arrival destination.³⁴ For team sports, the method, mode, distance, and timing of travel vary greatly and are largely dependent on scheduling, team budget, and the coach's preference.³⁶ Many teams, particularly in America and Australia, endure 1-way short-haul domestic or international travel up to 6 hours before or after competition.^{19,37,38} In addition to sleep disturbances, traveling can result in detrimental health, impaired mood, dehydration, and loss of motivation, all of which can affect recovery.³⁴ Of further concern, it has been shown that baseball teams whose circadian rhythms are more synchronized to optimal performance times are more likely to be successful, indicating either a negative effect of travel and/or desynchronized body-clock functioning.¹⁹ However, it should be noted that these data do not actually outline any physical or perceptual response to the travel, limiting their implication in athlete recovery.

Empirical data describing the effect of short-haul air travel on sleep, performance, and the ensuing recovery in these situations are largely lacking. For instance, the sleep quantity and quality of players after away-competition performance remain unclear, with short-haul air travel (1–3 h) affecting perceived sleep quality,³⁷ whereas some soccer players report earlier mean bedtimes after short-haul air travel (~5 h) and an away match.¹⁶ Competition performance, along with reduced physical demands, appears to be greater at home than away (in American football,³⁸ baseball,¹⁹ rugby league,¹⁴ and soccer¹⁶), suggesting either a negative effect of travel or a circadian advantage.³⁵ However, extrapolating these effects to determinations of match performance is difficult due to other external factors, the intermatch variability in opposition, and match intensity. While there have been few empirical studies, the available data suggest that short-haul travel has minimal effect on physiological and perceptual recovery (eg, no significant effect on YoYo Intermittent Recovery level 1 test performance), with more regular or longer periods of travel (eg, 24-h international transfers) more likely to result in negative responses.¹⁵ While short-haul air travel appears to have negligible effects on postmatch physiological recovery, the effect on perceptual markers of fatigue and sleep patterns after competition performance is equivocal. If these parameters decline, they can negatively influence training intensity or volume during ensuing sessions due to decreased motivation.³⁹ Given the myriad of conflicting demands while experiencing travel and sleep loss (eg, treatment, timing of training, recovery practices), it can be difficult for coaches to manage the most appropriate schedule for their team the day after a match. Indeed, more research is required to clarify the acute and chronic effects of cumulative travel (eg, over

a season) on sleep and psychological and physiological recovery parameters of professional team-sport athletes.

Sleep and Congested Competition Schedules

Excessive exercise loads can disturb the stress-recovery balance and result in performance decrements and injury occurrence.² For example, during periods of heavy match congestion in soccer, there is an increased injury risk for players when they play 2 matches per week rather than 1.⁴⁰ In this regard, some major European football teams may compete in up to 4 competitions at once—which likely affects players' sleep behavior. Congested schedules are also present throughout American sports such as baseball, hockey, and basketball. During these periods of high physical workloads, there is a potential for a reduction in sleep duration and quality. For example, it has been shown that as the effects of increased baseball match exposure accumulate toward the end of the season, strike-zone judgment is impaired, which suggests a fatigue-induced decline in performance, with sleep believed to be one of the main factors responsible.⁴¹

Sleep has also been suggested to be sensitive to exercise overload—with high training volumes associated with greater sleep disruptions.⁴² Although no published data are yet available in team-sport cases, Netzer et al⁴³ found significant increases in the REM-sleep-onset latency and decreases in REM sleep of well-trained cyclists after training and a competitive 120- to 150-km race, compared with no training or competition. Following this, it is logical that when team-sport athletes compete in a greater number of matches within a short period, exercise-induced muscle damage will accumulate (dependent also on exercise intensity), characterized by decreased neuromuscular function, increased perceptual fatigue, and increases in perceived soreness that can disrupt sleep.¹ Moreover, if there are several events in short succession, the continual anticipation of competition can also negate sleep.¹⁸ However, at present, there is little research that describes or quantifies the effect of these changes on subsequent recovery, particularly in team sports undertaking congested fixture scheduling. Future investigations into the time course of recovery after sleep loss would be particularly pertinent to team sports such as baseball and cricket, since these athletes can play on consecutive days and could be at a high risk of cognitive impairments (eg, slowed reaction time).

Sleep and Disturbances to Training Adaptation

Since sleep loss impedes muscle protein accumulation, the ability of skeletal muscle to adapt and repair can be hindered—which likely limits training adaptations.^{3,6,44} This may be concerning during the preseason for team-sport athletes given that sleep disturbances are present during higher training volumes.⁴² Since sleep loss can also affect vigor, mood, and perceptual awareness,³⁹ early training sessions could cause reductions in motivation and consequently reduce optimal training performance and subsequent adaptations.⁴⁵ Furthermore, if the stress-recovery balance of team-sport athletes is disrupted by either an increase in training load/stress or inadequate recovery, it may lead to an overreached, or even overtrained state.² Notably, disturbed sleep is believed to be one of many symptoms of either overreaching or the overtraining syndrome.² In a recent study, Hausswirth et al⁴⁶ found that objective measures of sleep duration and efficiency and immobile time were all negatively altered in a group of functionally overreached triathletes. There was also a higher prevalence of upper respiratory tract infections in this group, implying an association between the 2; however, whether impaired

sleep and illness occurrence are consequences, or simply symptoms or coincidental associations, of overreaching remains unknown.⁴⁶ In light of this, practitioners are encouraged to monitor the sleeping patterns of their athletes in high periods of stress either through subjective sleep diaries or wristwatch actigraphy.⁵

Since sleep loss can hinder the learning of new skills, affect emotional regulation, and disrupt cognitive function,⁶ it is likely that sleep is also important for optimizing cognitive training adaptations in team-sport athletes. For instance, sleep is critical for memory retention and neural plasticity and has been shown to improve visual discrimination and motor adaptation.²³ Therefore, it is likely that disturbing sleep during intense training or skill-acquisition periods (eg, preseason) will encumber adaptation in skill-based tasks with high neurocognitive reliance.⁴ However, objective evidence to support this suggestion is not currently available. Therefore, future research (with well-controlled randomized control trials) into the effects of sleep disruption on acute or chronic cognitive-based training adaptations in athletic populations is required.

Sleep Strategies for Team-Sport Athletes

Napping

In an attempt to recover from sleep debt, a commonly used sleep strategy among team-sport athletes is the restorative nap. Naps have been shown to improve alertness, sleepiness, short-term memory, and accuracy during reaction-time tests.⁴⁷ Furthermore, Waterhouse et al⁴⁷ found improvements in mean sprint performance after a 30-minute postlunch nap after 4 to 5 hours of sleep restriction. On the basis of this, it has been proposed that athletes take a postlunch nap to ameliorate the performance deficits caused by ultradian biological rhythms that occur within the circadian cycle.^{39,47} As such, it appears that napping behaviors have many benefits and should be undertaken where necessary in team-sport environments. An example would be for soccer players to have a nap after lunch if they are playing a match at night. However, it is critical that if naps are implemented in a team-sport environment they balance the need to enhance performance while not disturbing subsequent sleep patterns, as this could hinder the recovery process after training or competition. Indeed, while napping appears advantageous for performance (eg, napping before competition), more research is required to evaluate its possible effectiveness in recovery.

Sleep Extension

Extending sleep during normal sleep times is another strategy to alleviate the decrements in physiological and cognitive performance caused by sleep loss. Mah et al²⁴ found faster sprint and reaction times and improved shooting accuracy, energy, and mood after approximately 3 weeks of sleep extension (mean + 110 min) in 11 basketball players, indicating its use as a viable option for enhancing team-sport performance. Moreover, extending sleep improves psychological well-being, thus optimizing athletes' mental preparedness for competition.²⁴ However, obtaining extra sleep can be difficult, because increased sleep-onset latency and mood effects can be nullified due to earlier bedtimes. Thus, if an athlete is not sleep deprived it is possible that extending sleep will yield no benefit. The timing of this sleep intervention could also influence the effects of sleep extension, depending on the sleep chronotype of the athlete. In addition, more research assessing whether sleep extension during periods of high training load is a useful tool to ensure appropriate recovery is required. Such research would be pertinent in assisting

players achieve higher sustained intensities in subsequent exercise bouts (ie, during preseason).

Sleep-Hygiene Protocols

Identifying and modifying the factors that contribute to improve sleep quality (improving sleep hygiene) in team-sport athletes can also assist in ameliorating the detrimental effect of sleep loss and potentially enhance recovery. Sleep-hygiene strategies have been shown to improve sleep quality and onset latency in university students and to reduce sleep irregularity in adolescents, although the effect of numerous components of sleep hygiene in normal sleepers is mixed.⁴⁸ From an athletic perspective, little is known about the interaction between these sleep-hygiene strategies and the recovery of exercise and psychological parameters. Preliminary evidence indicates that adhering to some of the previous sleep-hygiene recommendations improves sleep quantity, resulting in a reduction in perceived soreness and fatigue in elite tennis players.⁴⁹ Furthermore, regulating sleep-wake times helps synchronize the circadian timing system, improving sleep quality and quantity.⁵⁰ As precompetition worry and anxiety are evident in athletes,^{10,18} it may be of benefit to use self-confidence tools (ie, meditation) to manage anxiety and stress, as these correlate with improved sleep.⁵⁰ Identifying each individual's best sleep habits (eg, bed comfort) is also pertinent, as unfamiliar environments may reduce sleep quality.⁵⁰ Such recommendations are similar to those designed for team-sport athletes who endure constant travel.³⁴ It is well known that sleep onset is prolonged by noise, light, and extreme temperatures, with athletes reporting noise and light as the 2 most important factors in their sleep quality.¹⁰ Since the use of technology just before sleeping promotes afferent signals from the retina to the pineal gland, inhibiting the secretion of melatonin and delaying sleep onset, the avoidance of bedtime technology (and thus reducing arousal and physiological excitement) has been recommended to improve sleep onset.⁵⁰ As part of a healthy sleep protocol, several nutritional recommendations have also been proposed to assist with sleep onset. For instance, a recent review by Halson⁵ proposed that diets high in carbohydrates

and protein may result in shorter sleep latencies and improved sleep quality, respectively.⁵ While there is a clear need for nutrition during the postexercise recovery period, the interaction between foods consumed postexercise and the ensuing sleep and recovery timeline is unclear. Indeed, the effects of nutrition are intricately complex and beyond the scope of this review (see Halson⁵ for further detail).

Future Research

Currently, there is insufficient evidence to conclusively describe the role of sleep for postexercise recovery and resultant performance outcomes. As such, the first step in understanding this contribution is to undertake long-term observational field studies through the use of subjective sleep diaries and/or actimetry in various situations. This will help identify areas where sleep may be an issue in team-sport athletes. Once this specific context is known, it is important to understand the interaction sleep has with variables in the high-performance athletic environment during situations where sleep is an issue. This requires both randomized crossover trials that investigate the measurement of sleep and the postexercise recovery timeline (both physiological and psychological) and also case studies in high-performance team-sport athletes. Future work in this field could also focus on understanding the mechanisms involved and providing appropriate interventions to improve sleep and the ensuing recovery process.

Practical Recommendations for Team-Sport Athletes

The recommendations in Table 1 are based on the literature in this review. However, we recognize that there is a lack of research examining the interactions between sleep and recovery in athletes. Nonetheless, there seems little risk but much (potential) benefit in following these recommendations. It is perhaps most important to tailor interventions toward individual athletes.

Table 1 Practical Sleep Recommendations for Players, Coaches, and Practitioners

Issue	Response
Determine whether there are sleep problems during normal scenarios in your athletic population.	One can do this by using subjective sleep diaries or wristwatch actimetry. Treat it in conjunction with a trained medical professional. Accommodating morning and evening types in team sports would appear particularly difficult, thus warranting clear communication between players, medical staff, and coaches.
Late-night matches and congested schedules.	Conduct correct sleep-hygiene practice after competition. This includes no technology 30 min before bedtime, no TV or use of laptops in bed, and dark, cool (but not cold), quiet rooms (blinds closed). Set a regular sleep schedule where possible and introduce relaxation and meditation techniques if necessary. These will presumably affect each athlete differently due to the intraindividual variability in sleep requirement. For further detail the reader is directed to Halson ⁵ and Malone. ⁵⁰
Short-haul domestic or international travel.	When traveling, ensure adequate hydration and time meals appropriately (usually in synchronization with the arrival time zone), move around the transportation vessel where/when possible, and synchronize light exposure to the arrival time zone. For detailed recommendations see Samuels. ³⁴
It is important that teams be aware of the possible altered physiological load in next-day training sessions after sleep loss.	Given the association between sleep loss and injury, ³³ individualized training after periods of sleep loss would seem appropriate. In general, advise and remind athletes to achieve consistent and adequate sleep (7–10 h/night), especially after a match.
Daytime sleepiness.	Napping appears beneficial for both repaying sleep debt and benefiting acute performance outcomes. However, be conscious of the effect of naps as they may also compromise recovery by interfering with subsequent sleep patterns.

Conclusion

While sleep is commonly reported by athletes, coaches, and scientists to be critical for recovery from intense exercise and/or competition, the current understanding of the effect of sleep on the recovery profile, especially in athletic populations, remains unclear. There is evidence to suggest that elite athletes lose sleep before and during competition periods. Furthermore, although limited published data are available, team-sport athletes appear to be susceptible to reductions in sleep quality and duration during and after competition (especially at night) and during periods of congested fixture scheduling and longer forms of travel. Given the regularity with which numerous professional teams might encounter these situations throughout a season, they may encumber the players' sleep and recovery. The efficacy of interventions to improve sleep, such as sleep-hygiene protocols and sleep extension, appears advantageous—but requires further investigation in situations relevant to professional team sports. These interventions may be suited to specific situations when the risk of compromised sleep is higher (ie, playing at home or away, at night, and/or inclusive of travel). This is especially pertinent with regard to the recovery of exercise parameters. Indeed, since research in this area is lacking, further research into the role of sleep and recovery in team sports is warranted.

Acknowledgments

Hugh Fullagar is supported by a Science and Health in Soccer scholarship funded by the DAAD (German Academic Exchange Service).

References

- Nédélec M, McCall A, Carling C, et al. Recovery in soccer: part II—recovery strategies. *Sports Med*. 2013;43(1):9–22. [PubMed doi:10.1007/s40279-012-0002-0](#)
- Kellmann M. Preventing overtraining in athletes in high-intensity sports and stress/recovery monitoring. *Scand J Med Sci Sports*. 2010;20:95–102. [PubMed doi:10.1111/j.1600-0838.2010.01192.x](#)
- Halsen SL. Nutrition, sleep and recovery. *Eur J Sport Sci*. 2008;8(2):119–126. [doi:10.1080/17461390801954794](#)
- Fullagar HHK, Skorski S, Duffield R, et al. Sleep and athletic performance: the effects of sleep loss on exercise performance, and physiological and cognitive responses to exercise. *Sports Med*. 2015;45(2):161–186. [PubMed](#)
- Halsen SL. Sleep in elite athletes and nutritional interventions to enhance sleep. *Sports Med*. 2014;44(Suppl 1):S13–S23. [PubMed doi:10.1007/s40279-014-0147-0](#)
- Samuels C. Sleep, recovery, and performance: the new frontier in high-performance athletics. *Neurol Clin*. 2008;26(1):169–180. [PubMed doi:10.1016/j.ncl.2007.11.012](#)
- Krystal AD, Edinger JD. Measuring sleep quality. *Sleep Med*. 2008;9(Suppl 1):S10–S17. [PubMed doi:10.1016/S1389-9457\(08\)70011-X](#)
- Skein M, Duffield R, Minett G, et al. The effect of overnight sleep deprivation after competitive rugby league matches on postmatch physiological and perceptual recovery. *Int J Sports Physiol Perform*. 2013;8:556–564. [PubMed](#)
- Erlacher D, Ehrlenspiel F, Adegbesan OA, et al. Sleep habits in German athletes before important competitions or games. *J Sports Sci*. 2011;29(8):859–866. [PubMed doi:10.1080/02640414.2011.565782](#)
- Venter RE. Perceptions of team athletes on the importance of recovery modalities. *Eur J Sport Sci*. 2014;14:S69–S76. [PubMed doi:10.1080/17461391.2011.643924](#)
- Lastella M, Lovell GP, Sargent C. Athletes' precompetitive sleep behaviour and its relationship with subsequent precompetitive mood and performance. *Eur J Sport Sci*. 2014;14(Suppl 1):S123–S130. [PubMed](#)
- Lastella M, Roach G, Halsen SL, et al. Sleep/wake behaviour of endurance cyclists before and during competition. *J Sports Sci*. 2015;33(3):293–299. [PubMed](#)
- Lastella M, Roach G, Halsen SL, et al. Sleep/wake behaviours of elite athletes from individual and team sports. *Eur J Sport Sci*. 2015;15(2):94–100. [PubMed](#)
- McGuckin TA, Sinclair WH, Sealey RM, et al. The effects of air travel on performance measures of elite Australian rugby league players. *Eur J Sport Sci*. 2014;14:S116–S122. [PubMed doi:10.1080/17461391.2011.654270](#)
- Fowler P, Duffield R, Vaile J. Effects of simulated domestic and international air travel on sleep, performance, and recovery for team sports. *Scand J Med Sci Sports*. 2015;25(3):441–451. [PubMed](#)
- Fowler P, Duffield R, Vaile J. Effects of domestic air travel on technical and tactical performance and recovery in soccer. *Int J Sports Physiol Perform*. 2014;9(3):378–386. [PubMed doi:10.1123/IJSP.2013-0484](#)
- Eagles A, McLellan C, Hing W, et al. Changes in sleep quantity and efficiency in professional rugby union players during home based training and match-play [Published online ahead of print November 4, 2014]. *J Sports Med Phys Fitness*. [PubMed](#)
- Juliff LE, Halsen SL, Peiffer JJ. Understanding sleep disturbance in athletes prior to important competitions. *J Sci Med Sport*. 2015;18(1):13–18. [PubMed](#)
- Winter WC, Hammond WR, Green NH, et al. Measuring circadian advantage in Major League Baseball: a 10-year retrospective study. *Int J Sports Physiol Perform*. 2009;4(3):394–401. [PubMed](#)
- Meyer T, Wegmann M, Poppendieck W, et al. Regenerative interventions in professional football. *Sports Orthop Traumatol*. 2014;30:112–118.
- Bruckner P, Khan K. *Clinical Sports Medicine*. Sydney, Australia: McGraw Hill; 2006.
- Vyazovskiy VV, Delogu A. NREM and REM sleep: complementary roles in recovery after wakefulness. *Neuroscientist*. 2014;20(3):203–219. [PubMed doi:10.1177/1073858413518152](#)
- Stickgold R. Sleep-dependent memory consolidation. *Nature*. 2005;437:1272–1278. [PubMed doi:10.1038/nature04286](#)
- Mah CD, Mah KE, Kezirian EJ, et al. The effects of sleep extension on the athletic performance of collegiate basketball players. *Sleep*. 2011;34(7):943–950. [PubMed](#)
- Schaal K, Le Meur Y, Louis J, et al. Whole-body cryostimulation limits overreaching in elite synchronized swimmers. *Med Sci Sports Exerc*. 2015;47(7):1416–1425. [PubMed](#)
- McMurray RG, Brown CF. The effect of sleep loss on high intensity exercise and recovery. *Aviat Space Environ Med*. 1984;55(11):1031–1035. [PubMed](#)
- Robey E, Dawson B, Halsen S, et al. Effect of evening postexercise cold water immersion on subsequent sleep. *Med Sci Sports Exerc*. 2013;45(7):1394–1402. [PubMed doi:10.1249/MSS.0b013e318287f321](#)
- Souissi N, Chtourou H, Aloui A, et al. Effects of time-of-day and partial sleep deprivation on short term maximal performances of judo competitors. *J Strength Cond Res*. 2013;27(9):2473–2480. [PubMed doi:10.1519/JSC.0b013e31827f4792](#)
- Myllymäki T, Kyröläinen H, Savolainen K, et al. Effects of vigorous late-night exercise on sleep quality and cardiac autonomic activity.

- J Sleep Res.* 2011;20(1 Pt 2):146–153. [PubMed doi:10.1111/j.1365-2869.2010.00874.x](#)
30. Raymond I, Nielsen TA, Lavigne G. Quality of sleep and its daily relationship to pain intensity in hospitalized adult burn patients. *Pain.* 2001;92:381–388. [PubMed doi:10.1016/S0304-3959\(01\)00282-2](#)
 31. Roach GD, Schmidt WF, Aughey RJ, et al. The sleep of elite athletes at sea level and high altitude: a comparison of sea-level natives and high-altitude natives (ISA3600). *Br J Sports Med.* 2013;47:i114–i120. [PubMed doi:10.1136/bjsports-2013-092843](#)
 32. Robey E, Dawson B, Halson S, et al. Sleep quantity and quality in elite youth soccer players: a pilot study. *Eur J Sport Sci.* 2014;14(5):410–417. [PubMed](#)
 33. Luke A, Lazaro RM, Bergeron MF, et al. Sports-related injuries in youth athletes: is overscheduling a risk factor? *Clin J Sport Med.* 2011;21(4):307–314. doi:10.1097/JSM.0b013e3182218f71
 34. Samuels CH. Jet lag and travel fatigue: a comprehensive management plan for sport medicine physicians and high-performance support teams. *Clin J Sport Med.* 2012;22:268–273. [PubMed doi:10.1097/JSM.0b013e31824d2eeb](#)
 35. Reilly T. How can travelling athletes deal with jet-lag? *Kinesiology.* 2009;41:128–135.
 36. Bishop D. The effects of travel on team performance in the Australian national netball competition. *J Sci Med Sport.* 2004;7(1):118–122. [PubMed doi:10.1016/S1440-2440\(04\)80050-1](#)
 37. Richmond LK, Dawson B, Stewart G, et al. The effect of interstate travel on the sleep patterns and performance of elite Australian Rules footballers. *J Sci Med Sport.* 2007;10(4):252–258. [PubMed doi:10.1016/j.jsams.2007.03.002](#)
 38. Smith RS, Guilleminault C, Efron B. Circadian rhythms and enhanced athletic performance in the National Football League. *Sleep.* 1997;20(5):362–365. [PubMed](#)
 39. Reilly T, Edwards B. Altered sleep–wake cycles and physical performance in athletes. *Physiol Behav.* 2007;90(2-3):274–284. [PubMed doi:10.1016/j.physbeh.2006.09.017](#)
 40. Dupont G, Nedelec M, McCall A, et al. Effect of 2 soccer matches in a week on physical performance and injury rate. *Am J Sports Med.* 2010;38(9):1752–1758. [PubMed doi:10.1177/0363546510361236](#)
 41. Kutscher S, Song Y, Wang L, et al. Validation of a statistical model predicting possible fatigue effect in Major League Baseball. *Sleep.* 2013;36:A408.
 42. Taylor SR, Rogers GG, Driver HS. Effects of training volume on sleep, psychological, and selected physiological profiles of elite female swimmers. *Med Sci Sports Exerc.* 1997;29(5):688–693. [PubMed doi:10.1097/00005768-199705000-00016](#)
 43. Netzer NC, Kristo D, Steinle H, et al. REM sleep and catecholamine excretion: a study in elite athletes. *Eur J Appl Physiol.* 2001;84:521–526. [PubMed doi:10.1007/s004210100383](#)
 44. Dattilo M, Antunes HKM, Medeiros A, et al. Sleep and muscle recovery: endocrinological and molecular basis for a new and promising hypothesis. *Med Hypotheses.* 2011;77(2):220–222. [PubMed doi:10.1016/j.mehy.2011.04.017](#)
 45. Sargent C, Halson S, Roach GD. Sleep or swim?: early-morning training severely restricts the amount of sleep obtained by elite swimmers. *Eur J Sport Sci.* 2014;14:S310–S315. [PubMed doi:10.1080/17461391.2012.696711](#)
 46. Hausswirth C, Louis J, Aubry A, et al. Evidence of disturbed sleep and increased illness in overreached endurance athletes. *Med Sci Sports Exerc.* 2014;46(5):1036–1045. [PubMed doi:10.1249/MSS.0000000000000177](#)
 47. Waterhouse J, Atkinson G, Edwards B, et al. The role of a short post-lunch nap in improving cognitive, motor, and sprint performance in participants with partial sleep deprivation. *J Sports Sci.* 2007;25(14):1557–1566. [PubMed doi:10.1080/02640410701244983](#)
 48. Stepanski EJ, Wyatt J. Use of sleep hygiene in the treatment of insomnia. *Sleep Med Rev.* 2003;7(3):215–225. [PubMed doi:10.1053/smr.2001.0246](#)
 49. Duffield R, Murphy A, Kellett A, et al. Recovery from repeated on-court tennis sessions: combining cold-water immersion, compression, and sleep interventions. *Int J Sports Physiol Perform.* 2014;9(2):273–282. <http://dx.doi.org/10.1123/IJSP.2012-0359>
 50. Malone SK. Early to bed, early to rise?: an exploration of adolescent sleep hygiene practices. *J Sch Nurs.* 2011;27(5):348–354. [PubMed doi:10.1177/1059840511410434](#)